

420

The
Manual of
Homœopathic Injections
&
Surgery.

(COMBINED)

BY

J. M. MITTER, B. A., M. D. (U. S. A.)

Principal, Modern Homœo.

Medical College etc.

MODERN ECONOMIC PHARMACY,
83, CLIVE STREET,
CALCUTTA.



420



The Manual of Homœopathic Injections & Surgery.

(COMBINED.)

BY

J. M. Mitter. B. A., M. D. (U. S. A.)

PRINCIPAL, THE MODERN HOMŒO. MEDICAL COLLEGE,

AUTHOR, MODERN MATERIA MEDICA, DIARRHŒA, FEVER ETC

EDITOR, "Homœopathist" & "Chikitsa-Samachar"

THE CHIEF RESEARCH-WORKER,

THE RESEARCH HOMŒO. ASSOCIATION OF INDIA,

VICE-PRESIDENT,

ALL BENGAL HOMŒO. CONFERENCE (1931),

ASST. SECRETARY, INSTITUTE OF HOMŒOPATHY

LATE DIRECTOR, THE RESEARCH HOMŒO. SOCIETY LD. ETC.



Price Re. ~~1~~/8 only.

Published by Dr. J. M. Mitter
13, Kailash Bose's Street, Calcutta.

HOMEO. INJECTIONS
OF
New Remedies

BY
J. M. Mitter, B. A., M. D. (U. S. A)

It contains therapeutic indications of infalliable injections for Asthma (Neo-Adrenaline), Brain-fag, Bubo, Birth control, Catarract, Carbuncles, Caries of bones, Eczema, Lumbago, Plague, Pneumonia, Piles, Ulcers, Leucoderma, Orchitis & Hydrocele, Rachitis, Uterine diseases, Vetigo, full treatment of Typhoid etc.

Price Rupee 1/- one only.

PRINTED BY
BEPIN BEHARY MONDAL
UNITED PRESS,
29, Grey Street, Calcutta.

INTRODUCTION,

Uptill now the Homœopathic Medicines were generally administered through the mouth or by olfaction only. But it has been found and declared by the eminent Homœopathists such as, Drs. G. Royal, Bœricke etc. that the Homœopathic Medicines often bring the best and quickest results when applied by hypodermic injections. But sometimes it is asked by the Homœopathists who do not like this innovation, wæther the process is in accordance with the doctrine of Hahnemann.

If with an unbiased mind we read the writings of Hahnemann, we would find that the theory of Homœopathic injection is not only not against the teachings of Hahnemann but is supported by his writings.

In the *Chronic Diseases* (Page 265) the founder of Homœopathy says, "But if the Diseased Organism is affected by the Physician through this same appropriate remedy at the same time in some *sensitive spots* other than the nerves of the mouth and alimentary canal *i.e.* if this same remedy that has been found useful is at the same time in its watery solution *rubbed* in (even in small quantities) into one or *more parts of the body which are free* from the morbid ailments (*e. g.* on an arm or on the thigh, leg, which have neither cutaneous eruption, nor pains

nor cramp) then the curative effects are much increased." In another place the Master writes, "When the small-pox or the cow-pox catches, this happens in the moment when in vaccination the morbid fluid in the bloody scratch of the skin comes in contact with the *exposed nerve*, which then, irrecoverably, dynamically communicate the disease to the Vital force (to the whole nervous system) in the same moment." Page 72. *The Chronic Diseases*.

As the Homœopathic Medicines act through the sentient nerves, a physician's duty is to bring the selected medicine in contact those nerves. This can be easily done by injecting the medicines under the skin of the body.

Dr. G. Royal writes that in Typhoid Fever where all the mucous membrane of the alimentary canal became ulcerated, it is the best plan to administer the indicated medicines by sub-cutaneous injections.

We do not decry the orthodox mode of administration of medicines through the mouth or by olfaction. All that we want to impress on the mind of the unbiased Homœopathists is that the administration of Homœopathic medicines by injections is not at all against the doctrine as expounded by Hahnemann ; it is only another scientific and effective weapon added in armoury of Homœopathy.



The Manual of Homœopathic Injections and Surgery. (COMBINED.)

Injection.

GENERAL HINTS.

Use strict antiseptic precautions. Cleanse the Hypodermic Syringe with absolute alcohol ; boil and cleanse it again before using.

Location of operation should be thoroughly rubbed with a piece of absorbent cotton saturated with absolute alcohol. **Inject Slowly.** On withdrawal of the needle after the injection the puncture should be pressed and rubbed with a thumb for a few minutes to prevent the oozing out of the medicine.

Ampoules—Each ampoule contains sufficient liquid for one injection, so no addition of extra liquid is generally necessary.

Syringe—Two C. C. Hypodermic glass Syringe will generally serve the purpose.

The Operator's hands—Wash the hands with a Soap and then dry them on a sterile towel and rub absolute alcohol on them thoroughly.

Mode of Injection—All injections excepting **Maloquin** No. 1, should be given hypodermically. On breaking the elongated top portion of the ampoule the needle of the syringe is to be inserted into its cavity to draw up its contents into syringe.

Location—Arm, thigh or buttock.

Caution—Do not injection on the affected part.

N. B. In case of inflammation of the part of the body where injection would be operated, foment it with a hot small cloth bag filled with ordinary salt.

ANTI-ABORTION.

Uses—In case of frequent and early miscarriage causing seemingly sterility it will save the fœtus.

Dose—1. c. c. once a week for at least two months. During the pregnancy Asokine may be used with great benefit.

General Measure—Instruct to avoid sexual congress during the pregnancy.

ANTI-TOXIN.

Uses—It is an excellent Small Pox prophylactic & curative.

Dose—1. c. c. every third day. Stop when the patient feels feverish. As a curative use 1. c. c. every day.

ADRENALINE.

It is the active principle of the medulla of the suprarenal gland.

Uses—It is invaluable in checking capillary hemorrhages from all parts of the body. It is useful in palliating acute attack of Asthma (for its permanent cure the **Pulmotone** is the best. Its indications and Dosage will be found in my “**New Remedies**”).

It temporarily relieves Angina Pectoris, Pericarditis and other heart troubles. It overcomes shock or heart failure during anesthesia.

Caution—The ampoules should be protected from light.

Dose—1. c. c. In urgent cases it may be repeated every 3 or 4 hours. It must not be too frequently repeated.

ACCELERATINE.

Uses—For general debility.

Indication—Nervous exhaustion, sexual debility. Nerve tremors of the aged ; Chorea ; Paralysis agitans ; Insomnia.

Dose.—1. c. c. Once a week.

ANÆMINE.

Uses—In pernicious Anæmia it is almost a specific.

Indications—Breathlessness, dizziness, headache, faintness, palpitation, fatigue, muscle cramp and cynosis ; also drawsiness, depression and insomnia.

Dose—1. c. c. Once every 3rd day until improvement begins.

ANTI-ZOOTAN.

No. 1 & 2.

Use—For scorpion bites, stings of poisonous insects and snake bites.

Indications—The patient feels in the bitten finger or toe an intense hot burning pain which has the tendency to ascend the arm or leg and a sensation of numbness and slowness of movement of the parts which likewise has a tendency to ascend and invade the whole limb. This sensation of numbness or torpor, rather soon, in an hour or so or in a few minutes in the most serious cases, changes to a paralysis, and even to a real spasmodic contracture of the injured limb which is very painful and alarming.

No. 1. Injection—1. c. c. used immediately after the stinging, ameliorates the burning pain produced and many times stops the progress of the poison.

No. 2. Dose—1. c. c. Every three hours. Should be used when the above mentioned symptoms fully develop.

ANTI-RHEUMATE.

No. 1.

Uses—For Rheumatism, acute and chronic. Gout and articular rheumatism, muscular pains and soreness. Pain worse, motion, better by pressure. Right side mostly affected. Chronic gout and stiffness of joints.

Dose—1. c. c. Every 3rd or 5th day-

No. 2.

Uses—In Rheumatism of females with uterine troubles.

Indications—The pains are wondering ; rheumatism of muscles of the limbs and trunk ; also articular Rheumatism of the lower extremities.

Dose—1. c. c. Once weekly.

ANTI-TUMOR.

No. 1.

Uses—Tumors, benign and malignant.

Indications—In hard, stony glands, varicose and enlarged Veins.

Dose—1. c. c. Every week.

No. 2.

Hard knots in female breasts.

Dose—1. c. c. Every week.

ANTI-BERI-BERI

Uses—For Beri-Beri and epidemic dropsy.

Dose—1. c. c. Every third day.

ANTI-INSANITY.

No. 1.

In acute mania.

Indications—Violent ; restless ; afraid of being poisoned ; delirious mania.

Dose—1. c. c. Every four hours.

No. 2.

Uses—For chronic mania.

Indications—Ill-humoured and quarrelsome ; delusions ; believes old rags to be silk. The head is hot and the feet cold or other evidence of disturbed circulation.

Dose—1. c. c. Once a week.

ANTI-TETANUS.

Uses—For Tetanus or Lockjaw.

No. 1.

Indications—Tetanus from wounds in tendons, sole of feet or of fingers and palm of hands.

Dose—1. c. c. Every three hours until improvement begins. With it Hypericum 200 may be administered with great advantage.

No. 2.

Indications—It is very useful when the tetanus is not due to any of the above-mentioned causes or no cause can be found but the following symptoms be present—there are spasms excited by the least cause, affecting first muscle of the jaw, neck and back, then the other muscles of the body. Dyspnœa is frequently present; pain through the body is violent; the mind is clear.

Dose :—First day 3. c. c. thrice; subsequent days gradually up to 3. c. c. four times a day.



CARDEIN.

As it is absolutely free from Digitalis and Strychine its use will leave no bad after-effect.

Uses—Useful for *chronic heart diseases*.

Indications--Chronic heart disease with extreme weakness. Very feeble and irregular action of heart. Cardiac dropsy. Extreme dyspnoea on least exertion, without much increase of pulse. Heart dilated; first sound weak. Pulse accelerated, irregular, feeble, intermittent.

Dose—1. c. c. Every fourth day or oftener according to the discretion of the attending physician.

CARISCURO.

Uses—Caries in bones.

No. 1.

Indications—Periostium painful, swollen and enlarged. Darting pain and caries in bones. Ulcers affecting bones. thin ichorous pus.

Dose—1. c. c. Every third day.

No. 2.

Indications—Caries of nasal, palatine and mastoid bones. Soreness of affected bones.

Dose—1. c. c. Once in a week.

CANCERADINE.

Uses—Epitheliomata, cancer of the mammæ, the uterus, the tongue or other parts or the body are cured by it.

Indications—Great pain and induration of mammary glands; offensive discharge and hæmorrhage of uterus with unbearable pain. Cancerus cachexia.

Dose—1. c. c. Every third day.

CHOLERATOX.

Uses—It is used as a prophylactic and a curative.

Indications—In cases of cholera morbus or the Asiatic Cholera when vomiting accompanies the purging.

Dose—1. c. c. As curative every one hour; and as prophylactic once a week

COLONUS.

USES.—For Leprosy, Elephantiasis, etc.

Indications—Tubercular or syphilitic leprosy; lassitude, indisposition to move; loss of energy; apathy and obstruction of the capillaries; intolerable itching over the whole body.

Dose—1. c. c. Every 2nd day.

About 9 injections are sufficient to cure the ulcers.

DELIVERINE.

Uses—(a) *For the safe delivery.*

If the medicine be used from the 7th month of pregnancy there will be no trouble in delivery.

Dose—1. c. c. every 15th day.

(b) *For quick delivery*—Inject 1. c. c. every two hours. It is also useful for **after-pains** with great sensitiveness and intolerance to pain.

Dose—1. c. c. Every two hours.

DIPHATHERINE.

Uses—Diphtheria ; malignant sore throat ; malignant quinsy.

Indications—Fever ; profound prostration ; soreness of the throat ; constant desire to hawk. The tonsils are swollen, upon which will be found the membrane. There is a *very offensive breath*, due to the diphtheric process. The glands of the neck are swollen and tender. The membrane may extend into the nose, called *nasal diphtheria*, and causes epistaxis, and a yellow offensive discharge from the nose, ulcerating the upper lip. If it should extend to the larynx (*laryngeal diphtheria*) it will cause hoarseness ; croupy cough ; dyspnœa and stridulous

breathing. If the membrane form upon the external mucous membrane it is called cutaneous diphtheria.

Dose—1. c. c. Every two hours.

Diet.—It is of first importance to keep the patient well nourished. Let the diet be in liquid form, highly nutritious, easily digestible. Milk is highly recommended ; must give in small quantities, at regular intervals, day and night ; if the patient can not swallow then give the food by enemata.

DOLVINA.

No. 1.

Uses—It is used with the best result in Amebic dysentery.

Indications—Little thirst ; while straining pain so great that it nauseates. Cutting, clutching ; wores, around the navel.

Dose—For adult 1. c. c. & for children above 3 years of age $\frac{1}{2}$ c. c. In acute case twice daily and in chronic case twice in a week.

No. 2.

Uses—In chronic Dysentery it gives satisfactory result.

Indications—When No. 1. does not prove to be very effective,

Dose—1. c. c. Every other day until improvement begins, then once a week for four weeks.

DIABATONA.

Uses—For Diabetes mellitus or Glycosuria.

No. 1.

Indications—Headache, malaise, unquenchable thirst and frequent urination. Despite the good appetite the patient constantly loses flesh.

The urine is greatly increased, pale in colour, a musty odor, the sp. gr. is high, from 1.025 to 1.050 and upon Chemical examination sugar will be found. *Great emaciation, debility and tendency to ascites and general dropsy.*

Dose—1. c. c. Every third day.

No. 2.

Indications—Urine like milk, mixed with jelly-like, bloody mucus or clear like water; pain in the back and region of the kidneys: Sleeplessness; excessive emaciation; great prostration. After loss of animal fluids; after grief, anguish, sorrow and care.

Dose—1. c. c. twice weekly.

DYSMENORINE.

Uses—For neuralgic, inflammatory, obstructive and membranous Dysmenorrhœa.

No. 1.

Indications—Menstruation too early, too profuse, offensive in odor, like jelly; or scanty, thin, light-coloured. Spasmodic, cramping pain, in lower abdomen and thighs, resembling labour pains.

Dose—1. c. c. Before or after the attack every third day.

N. B. During the attack use No. 3 1. c. c. every two hours.

No. 2.

Indications—Scanty flow; neuralgic and congestive dysmenorrhœa with spasmodic irregular and very severe pains, especially the first two days of menses; Pain in small of back and great aching and soreness of lower limbs, bad breath, bitter taste, vertigo, chilliness.

Dose—1. c. c. Before the attack and after the attack every third day. During the pain use No. 3. 1. c. c. every two or three hours.

No. 3.

It will palliate the pains without interfering with action of the main medicine.

Dose—1. c. c. Every two hours.

ECZONA.

Uses—Humid, spreading and scurfy eczema. Painful to contact, violent itching and burning, aggravation in the mornings, indoor, from cold or draught of air; at night with profuse serous exudates, gradually becoming sticky, gluey and finally offensive.

Dose—1. c. c. Every 10th day.

Note—For immediate relief of violent itching, burning etc. use "ECZONE OINTMENT."*

ERYSIPLINE.

Uses—For Erysipelas.

No. 1.

Indications—Oedematous erysipelas of the face, travel-

* Eczone Ointment can be had from all Pharmacies
Price 1/- per pot.

ing from right to left. Affected part assumes a pinkish rosy hue. There is stinging, burning, pricking pain. High fever with dry skin.

Dose—1. c. c. (in serious case) one in the morning and another in the evening.

Note—Cover the affected part with a piece of cloth saturated with normal Saline water.

No. 2.

Indications—Erysipelas covered with large blisters of smaller vessels covered with serum, beginning on the left side of face and spreading to the right. Affected part becomes dark-red.

Dose—1. c. c. in serious cases twice daily (morning and evening) and in milder cases one injection daily.

No. 3.

Indications—Facial erysipelas, especially left side, at first bright-red but it soon takes on a *dark bluish hue*.

Dose—1. c. c. One injection daily.

FILARIN.

Uses—For Elephantiasis and Filaria.

No. 1.

Indication—It is an infallible remedy of Filaria when

it aggravates with inflammatory high fever in new and full moon.

Dose—1. c. c. once in a week.

No. 2.

Indications—Great thickening of epidermoid layer and enfoliation of scales. It has a high reputation in leprosy and lupus, when there is no ulceration.

Dose—1. c. c. twice weekly.

FISTULUS.

Uses—For fistulæ in Ano.

No. 1.

Indications—It is almost a specific for fistula ano. Drs. Farington, Royal and Bœricke have highly recommended it. Old fistulous ulcers with a watery discharge mixed with pus which aggravates in full or new moon, are highly benefitted by this injection.

Dose—1. c. c. Every fort-night.

No. 2.

Indications—Fistula anus with great constriction, burning like fire which is temporarily relieved by cold water. Itching of anus.

Dose—1. c. c. Every 7th day.

No. 3.

Indications—Habitual constipation, patient often suffers from eczema. No tenesmus or constriction of anus.

Dose—1. c. c. Every seventh day.

GONORINUM.

Uses—For Gonorrhœa acute and chronic stages.

No. 1.

Indications—Acute gonorrhœa. Frequent urination with a burning pain ; acute gonorrhœa, aggravates in the evening ; urine flows slowly, has to force out the last drops ; great urging to urinate, with much straining, but cannot pass a drop. There is scalding and burning in the urethra before and after urination.

Violent and painful erections of male seminal organs, worse when walking and sitting, better when lying down. Sexual desire excessively increased. Acute inflammatory gonorrhœa with a copious, yellowish white discharge ; mucous discharge often glossy.

Dose—1. c. c. Every third day.

No. 3.

Indications—Burning at the meatus or at the orifice—a burning as of fire. Urine hot as it passes the orifice and burning continues for sometime after urination. Burning, cutting, scalding with tenesmus and chordee after urinating. Urine reddish or even bloody.

A thick creamlike discharge with a slight, greenish tinge to it.

Dose—1. c. c. Every 2nd day.

No. 4.

Indications—Gonorrhœa with stranguary and irresistible desire to urinate with crawling in the navicular fossa. Acute cystitis following the use of medicated injections in Gonorrhœa. Discharge is milky, albuminous, and sometimes yellow.

Dose—1. c. c. Every week.

No. 5.

Indications—Thick, yellow, purulent discharge and urethra feels sore and swollen, the last drops of urine seem to be retained within the urethra.

Dose—1. c. c. Every week.

No. 6.

Indications—Do not think of it in acute gonorrhœa. Loss of power to expel urine ; sensation as if urine or some

fluid were passing from urethra when not urinating. Sense of constriction of urethra, wants to pass urine but can not. Sub-acute or chronic Gonorrhœa with thin and greenish discharge. It is very effective as an inter-current medicine in the treatment of Gonorrhœa.

Dose—1. c. c. Every seventh day.

KALZINA.

Uses—For Kala-azar.

Indications—Enlarged and hard spleen and liver, cachexia. Lustreless hairs. **Darkening of the skin** which is more marked in the face. A tendency of bleeding from the gums and nose.

Dose—1. c. c. Every third day.

LEUCORUS.

Uses—For Leucorrhœa.

No. 1.

Indications—Painless leucorrhœa, discharge of thick, white mucous, milky, aggravates on lying down, burning, thin, arid, milky, mucous; thick and white leucorrhœa

before and during menses, with cutting pains in abdomen and cramps in bowels.

Dose—1. c. c. Every alternate day.

No. 2.

Indications—Leucorrhœa just midway between the menstrual period; albuminous white discharge, like white of eggs, or starch, with sensation as if warm water were escaping, perfectly bland and painless; great general nervousness.

Dose—1. c. c. Every third day.

No. 3.

Indications—Suitable to debilitated women. Leucorrhœa yellow greenish; with much stitches in uterus and great itching in Vagina and Vulva. Leucorrhœa aggravate after coition. Great bearing-down sensation. The patient is averse to household occupation, irritable and indifferent to those loved best.

Dose—1. c. c. Every seventh day.

MERC SALVINUS.

Uses—For Syphilis.

No. 1.

Indications—Red chancre on prepuce; spreading and deeply penetrating ulcer on glans and prepuce;

pale-red vesicles forming small ulcers after breaking, on glans and prepuce; painful bleeding chancres, with yellowish, foetid discharge; ulcers of glans and prepuce, with cheesy, lardaceous bottom and hard edges.

Dose—1. c. c. Every seventh day.

No. 2.

Indications—Generally useful in the 2nd stage of the disease and in old cases in scrofulous patients. Hunterian hard chancre; threatened gangrene of glans in paraphimosis; Soreness of bones of face; sharp shooting stiches in the end of the penis through the glans; old buboes, discharging for years.

Dose—1. c. c. Every week.

N. B. No. 1 & 2 Merc Salv may be alternately used in the 1st & 2nd stages of syphilis.

No. 3.

Indications—Secondary and tertiary syphilis, phagedenic chancres; Soreness of the skin and cranial bones; worse from damp weather; syphilitic ulcers in the throat; Copper-coloured spots over the body. Bad effects of the misuse of mercury.

Dose—1. c. c. Every fourth day.

No. 4.

Indications—Mental states of great depression; great

desire to commit suicide. Exostosis, Caries, nightly bone-pains, specially cranial, nasal and palative.

Dose—1. c. c. Every seventh day.

No. 5.

Indications—Chanere with hard edges and curly pus; *deep-eating ulcers*; Violent headache, causing hard lumps in head; indolent swollen glands. It is also useful in gummatous tumors involving the nervous tissues.

Dose—1. c. c. Every seventh day.

No. 6.

Indications - This remedy has achieved wonderful beneficial results after Mere salv. No. 5 failed to completely cure the syphilitic affection; Violent occipital headaches, dysphagia and syphilitic throat and bone affections; Caries of bones of feet.

Dose : -1, c. c. Every seventh day.

MALOQUIN.

Uses—For malarial fever.

No. 1.

Indications—Typical Malarial fever with three stages—the **chill**, the **heat** and the **sweat**. It is also

indicated in a case, where one or the other stages is wanting; and when order of its stages is reversed, so that, for example, the paroxysm commences with sweat and ends with the chill.

Dose—1. c. c. during the paroxysm and then every third day. To be given by **intermuscular** injection.

No 2.

Indications—Periodicity marked with adynamia.

Dose—1. c. c, Every third day, to be injected hypodermically.

NERVINOS.

Uses—For sexual weakness and impotency.

No. 1

Indications—Neurasthenic impotency. It is an excellent sexual stimulant. In spite of strong desire penis remains unerected.

Dose—1. c. c. Every 3rd day in the evening.

No. 2

Indications—Premature old age from abuse of sexual power. History of repeated gonorrhœa. No secretions.

Impotence. Parts cold, relaxed. *Desire gone.* Scanty emission without ejaculation.

Dose—1. c. c. Every 7th day.

NAT-CACO.

Uses—For anæmia and general debility.

Indications—Anæmia due to sudden loss of blood by hæmorrhage, excessive loss of semen, too long-continued lactation, profuse menstruation, albuminuria, diarrhœa, dysentery, malarial affection, diseases of spleen, and fever.

Dose—1. c. c. Every week.

PUR-CARBO.

Uses—For collapse in Pneumonia, Typhoid, Cholera, etc.

No. 1.

Indications—The patient may be almost life-less, but the head is hot; coldness, breath cool, pulse imperceptible, oppressed and quickened respiration and must have air, must be fanned hard, must have all the windows open.

Dose—1. c. c. Every three hours or earlier.

No. 2.

Uses—For collapses in cholera or after operation.

Indications :—Icy coldness of the whole body ; sudden sinking of strength ; pulse small and weak.

Dose—1. c. c. Every hour.

PYMONIN.

Use—For Septicæmia.

Indications—Dr. Allen says, “In septic fevers, especially puerperal, it has demonstrated its great value as a homœopathic dynamic antiseptic.” It is most effective in hectic, typhoid, typhus, ptomiane poisoning, dessecting wounds, after-effects of miscarriage. *All discharges are horribly offensive.* Great pain and burning in abscesses. Chronic complaints that date back to septic conditions.

Dose 1. c. c. Every 3rd day.

PHLOGIMUM.

Uses—For inflammations.

Dose—1. c. c. Every three hours.

PEDAPEPATINE.

Uses—For infantile liver.

Dose—1. c. c. Every seventh days.

Note—Don't allow the patient to take milk when the disease accompanies with fever. Goat's milk may be allowed when the fever subsides.

RADIARIN.

No. 1, 2.

Uses—For high blood pressure.

No 1.

Indications—Surging of blood to head and heart. Tendency to sudden and violent irregularities of circula-

tion, sensation of pulsation throughout the body. Can not bear any heat about the head. Vertigo on assuming upright position. Threatened apoplexy.

Dose—1. c. c. Every third day.

No. 2

Indications—Useful in chronic cases with same symptoms.

Dose—1. c. c. Every 7th day.

THEOGENIC.

No 1, 2 & 3.

Uses—For Phthisis.

No 1.

Indications—Generally used in the first stage of Phthisis. Hoarseness, racking cough and profuse expectoration of a virulent nature, attended with cardiac weakness, emaciation and general debility; diarrhœa.

Dose—1. c. c. Every 2nd day.

No, 2.

Indications—It is useful in the 2nd stage of the disease.

No. 3.

Indications—Generally used in the 3rd stage of Phthisis. Persistent inclination to cough. Cough attended with profuse greenish or mucopurulent expectoration; hectic

fever and emaciation ; coarse rales ; soreness in the chest after coughing ; sense of weakness in the chest ; talking causes fatigue ; expectoration sweetish in taste.

Dose—1. c. c. Every third or fourth day.

TYPHODINE.

Uses—For Typhoid fever.

Remarks—It produces a form of anti-bodies to the bac. Typhosus, viz., the agglutinius.

Indications—Great muscular soreness and putrid phenomena always present. All the secretions are offensive—breath, stool, urine, sweat etc.

Dose—1. c. c. daily.

N. B. It is advisable to inject **Pymonin** once a week as an intercurrent remedy,

ZINZI or ANTI-GANGRENE.

Uses—For gangrene, boils, carbuncles and ulcers.

Indications—Bluish, purple surroundings. The ulcers have an unhealthy discharges with offensive odour ; with the gradual spreading of ulcers lumps of dead flesh become disintegrated.

Dose—1. c. c. Every fourth day.





